

MASSIVE STARS IN THE SGR 1806–20 CLUSTER

DONALD F. FIGER¹, FRANCISCO NAJARRO², T. R. GEBALLE³, R. D. BLUM⁴, ROLF P. KUDRITZKI⁵

Draft version February 2, 2008

ABSTRACT

We report the discovery of additional hot and massive stars in the cluster surrounding the soft gamma repeater SGR 1806–20, based upon UKIRT and Keck near-infrared spectroscopy. Of the newly identified stars, three are Wolf-Rayet stars of types WC8, WN6, and WN7, and a fourth star is an OB supergiant. These three stars, along with four previously discovered, imply a cluster age of ~ 3.0 – 4.5 Myr, based on the presence of WC stars and the absence of red supergiants. Assuming coevality, this age suggests that the progenitor of SGR 1806–20 had an initial mass greater than $\sim 50 M_{\odot}$. This is consistent with the suggestion that SGRs are post-supernovae end states of massive progenitors, and may suggest that only massive stars evolve into magnetars that produce SGRs. It also suggests that very massive stars can evolve into neutron stars, not just black holes, as recently predicted by theory. The cluster age also provides constraints on the very high mass object, LBV 1806–20.

Subject headings: stars: evolution — stars: neutron — stars: individual (LBV1806-20) — stars: individual (SGR 1806–20) — infrared: stars

1. INTRODUCTION

Massive stellar clusters are the birth sites of the most massive stars, and are proving grounds for theories of massive star formation, evolution, end states, and Galactic chemical enrichment. With the recent advent of sensitive infrared instrumentation, on ground and space based platforms, many massive clusters in the Galaxy are now just being discovered and investigated. At least two of these clusters contain soft gamma repeaters (SGRs), a rare phenomenon characterized by persistent and energetic bursts of gamma ray emission lasting up to several seconds and releasing $\sim 10^{40}$ ergs s^{−1} (Mazets, Golenetskij, & Guryan 1979; Mazets et al. 1979; Kouveliotou et al. 1998). Only four SGRs are known, three in the Galaxy and one in the SMC. Three of the SGRs are associated with massive stellar clusters.

SGR 1806–20 is surrounded by a cluster of massive stars (Fuchs et al. 1999). Kulkarni et al. (1995) identified a luminous star that they claimed may be the infrared counterpart to SGR 1806–20, but Eikenberry et al. (2001) and Kaplan et al. (2002) determined that it is too far from the Chandra error box; no near-infrared counterpart has yet been found, even down to very faint magnitudes (Kouveliotou et al. 2004; Israel et al. 2004). van Kerkwijk, Kulkarni, Matthews, & Neugebauer (1995) find that this star has characteristics similar to those of Luminous Blue Variables (LBVs), i.e. it has $L \gtrsim 10^6 L_{\odot}$ and a spectral type in the range O9–B2. Eikenberry et al. (2004) estimated an extremely high

mass for LBV 1806–20, $\sim 200 M_{\odot}$; but Figer et al. (2004) have shown that it has double lines and thus may be binary.

In this Letter we present near-infrared spectra of fifteen bright sources near SGR 1806–20. We find that three of them are massive stars in the advanced Wolf-Rayet stage of post-main sequence evolution, and another three are less evolved OB supergiant stars. We use these results to constrain the nature of SGR 1806–20.

2. OBSERVATIONS AND DATA REDUCTION

Data were obtained at UKIRT⁶ on 7 and 8 June 2004, using UIST (Ramsay Howat et al. 1998) in medium resolution mode ($R \sim 800 = \lambda / \Delta \lambda_{\text{FWHM}}$, 0".24 slit width) from 1.50 μm to 2.50 μm . One or two 60 second exposures were obtained, after which the telescope was nodded along the direction of the slit and the similar exposures obtained. Star HIP86814 (F6V) was observed as a telluric standard; its Brackett series lines were removed by interpolation prior to ratioing.

Additional imaging and spectroscopic data were obtained at Keck⁷ on 22 June 2003, using NIRSPEC (McLean et al. 1998), with the NIRSPEC-6 filter, which covers the H and K bands). The slit-viewing camera (SCAM) exposures were 2 seconds and provided stellar images having FWHM $\sim 0".4$. Figure 1 shows a SCAM image; note the faint image produced by the slit. Spectra covering 1.6–2.35 μm were obtained at high resolution ($R \sim 26,500 = \lambda / \Delta \lambda_{\text{FWHM}}$, 0".45 slit width), using two 100 second exposures per grating setting, with the telescope nodded along the direction of the slit length in between exposures. Star #3 of the Quintuplet cluster was observed as a telluric standard (Figer et al. 1999).

⁶ The United Kingdom Infrared Telescope (UKIRT) is operated by the Joint Astronomy Centre on behalf of the Particle Physics and Astronomy Research Council.

⁷ Data presented herein were obtained at the W.M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W.M. Keck Foundation.

Electronic address: figer@stsci.edu

¹ Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218; figer@stsci.edu

² Instituto de Estructura de la Materia, CSIC, Serrano 121, 29006 Madrid, Spain

³ Gemini Observatory, 670 N. A'ohoku Pl., Hilo, HI 96720

⁴ Cerro Tololo Interamerican Observatory, Casilla 603, La Serena, Chile

⁵ Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822




FIG. 1.— NIRSPEC/SCAM image of stars in the 1806–20 cluster with coordinates offset from LBV 1806–20 at $18^h08^m40.312^s$ $-20^\circ24'41''14$ (J2000). Number designations are from this paper. Letter designations are from Eikenberry et al. (2004). Stars #8, 9, and 10 are to the west of the field. The slit can be seen as a linear artifact superpositioned on images of Star “C” and “B.” A circle marks the location of SGR 1806–20 (Kaplan et al. 2002).

For both sets of spectra arc lamps were observed to set the wavelength scale and continuum lamps provided “flat” images. Data were reduced by differencing the nod pair frames to remove background flux, dark current, and residual bias and dividing the result by the flat field frame in order to compensate for non-uniform response and illumination of the array. Finally, each extracted target spectrum was divided by the spectrum of a telluric standard.

Table 1 lists the coordinates, magnitudes, spectral types, and dates observed for the massive stars and field red giants. Figures 2, 3, and 4 show spectra of these stars.

3. DISCUSSION

3.1. Photometry and Spectral Analysis

Photometry was mostly obtained from the 2MASS⁸ (Skrutskie et al. 1997) Point Source Catalog (PSC). Stars #1, #3, and #4 are not in the catalog, so we estimated photometry for them from the Keck SCAM image using the 2MASS PSC to set a zero point. As a check, photometry for stars #7 and #11 from the Keck SCAM image differ from the PSC by +0.11 and −0.07 magnitudes, respectively. Note that our method relies on the fact that the NIRSPEC-6 filter transmission function, convolved with the spectral energy distributions with the appropriate extinction of the target stars, gives similar photometry as measured with the K filter used in 2MASS. Star “B” and “D” are in crowded regions, so we use photometry for them from Eikenberry et al. (2004),

⁸ This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

TABLE 1
MASSIVE STARS IN THE 1806–20 CLUSTER

ID	R.A.	Dec.	H–K	K	Type	Dates Observed
1	38.32	33.5	NA	11.76	WC8	7 June 2004
2	39.42	42.57	1.87	12.16	WN6	8 June 2004
3	39.50	35.88	NA	12.87	WN7?	8 June 2004
4	39.16	32.88	NA	12.45	OBI	22 June 2003
5	38.13	34.77	1.68	9.25	RG	7 June 2004
6	38.12	43.35	1.42	10.96	RG	7 June 2004
7	38.75	26.48	1.49	11.90	OBI?	7 June 2004
8	36.72	54.27	1.40	8.70	RG	8 June 2004
9	36.72	27.80	1.45	8.89	RG	8 June 2004
10	37.96	16.47	2.95	10.09	RG	8 June 2004
11	38.52	39.95	1.66	11.92	OBI?	8 June 2004
12	38.57	13.55	1.42	10.81	RG	8 June 2004
13	39.49	11.77	NA	11.16	NA	8 June 2004
14	39.40	13.02	NA	12.00	RG	8 June 2004
A	40.31	41.14	NA	9.26	LBV	22 June 2003
B	39.24	31.86	2.94	10.50	WC9	22 June 2003
C	39.51	30.02	1.85	10.96	OBI	22 June 2003
D	39.51	35.91	1.73	11.11	OBI	22 June 2003

NOTE. — Number designations are from this paper. Letter designations are from Eikenberry et al. (2004). Coordinates are offset from Right Ascension 18^h08^m and Declination $-20^\circ24'$ (J2000), and are from the 2MASS PSC, except for #3, #4, #13, and #14 which were estimated from the Keck SCAM image. Photometry are from the PSC, except for #1, #3, #4, #13, #14, and “A” which are estimated from the Keck SCAM image, and “B” and “D” which are from Eikenberry et al. (2004). The spectral type for “B” is taken from Eikenberry et al. (2004). Star #2 is source “E” in Eikenberry et al. (2001). The “RG” label refers to red giants.

instead of the PSC. As a further consistency check on the photometric systems used in the Table, note that 2MASS and Eikenberry et al. (2001) photometry match to within the quoted uncertainties of 0.1 magnitudes for Star #2.

We estimated spectral types by comparing the observed spectra to those in the Wolf-Rayet spectral atlas of Figer, McLean, & Najarro (1997) and Hanson, Conti, & Rieke (1996).

The spectra in Figure 2 reveal three new WR stars. Star #1 has strong lines of C III ($2.11\ \mu\text{m}$), C IV ($2.08\ \mu\text{m}$), He I ($1.70\ \mu\text{m}$ and $2.06\ \mu\text{m}$), and He II ($1.87\ \mu\text{m}$ and $2.16\ \mu\text{m}$). Taken together, these lines indicate a classification of WC8. Stars #2 and #3 have strong lines of He II; neither has appreciable N V ($2.10\ \mu\text{m}$) emission. Star #2 has relatively strong He II ($1.87\ \mu\text{m}$ and $2.16\ \mu\text{m}$), and $\text{EW}_{2.189\ \mu\text{m}}/\text{EW}_{2.112\ \mu\text{m}}=2.3$, consistent with a WN6 spectral type (Figer, McLean, & Najarro 1997). For star #3, we suggest a classification of WN7, but note that the spectrum has relatively low S/N.

The spectra in Figure 3 show three less evolved OBI stars (“4”, “C”, and “D”), as well as the evolved luminous blue variable, LBV 1806–20, and a previously identified WR star, “B” (Eikenberry et al. 2004). LBV 1806–20 is discussed at length in Figer et al. (2004), and references therein. The OBI stars show modest He I and H I absorption components with equivalent widths of $\text{EW}_{2.112\ \mu\text{m}}\sim 2\text{--}3\ \text{\AA}$ and $\text{EW}_{2.166\ \mu\text{m}}\sim 3\text{--}5\ \text{\AA}$, and $\text{FWHM}\sim 70\text{--}80\ \text{km s}^{-1}$; both line depths and equivalent widths are consistent with the classifications of O9–




FIG. 2.— UKIRT/UIST spectra of the newly discovered WR stars. The number labels refer to objects in Table 1.




FIG. 3.— Keck/NIRSPEC spectra of massive stars near the 2.112/2.113 μm He I doublet (left), and the 2.166 μm Br- γ line (right). The spectra (except for “B”) show the He I doublet in absorption near 2.1125 μm is marked by vertical lines. This feature in LBV 1806–20 is obviously double. The broad feature near 2.089 μm is common to LBV spectra and corresponds to an Fe II transition. Weak He I absorption lines just short of 2.166 μm are visible in all spectra, except for “B.”

B3 for these stars (Hanson, Conti, & Rieke 1996). Note that our high-resolution spectra successfully resolve these individual lines that would otherwise appear as blends in lower resolution spectra. Indeed, our linewidths are significantly less than those estimated in much lower spectral resolution data of Eikenberry et al. (2004), and they do not support the suggestion that these stars are extraordinarily luminous hypergiants.

Figure 4 demonstrates that most of the remaining sources are red giants (RG’s), given their strong CO absorptions at 2.3 μm and beyond. These stars are relatively old ($\tau_{\text{age}} > 100$ Myr) and have no physical association with the cluster. Stars #7 and #11 lack CO and thus could be OBI stars, but they lack Br- γ absorption




FIG. 4.— UKIRT/UIST spectra of stars within and near the field in Figure 1. Most of the spectra have CO bandhead absorption starting at 2.293 μm and characteristic of red giants.

lines. Pa- α absorption lines in their spectra would be comparable to the noise levels, so it is difficult to rule out their existence.

3.2. Massive Stars and SGR 1806–20

The 1806–20 cluster contains at least four WR stars, three OBI stars, and an LBV-candidate. While such a collection of spectral types may seem extraordinary, it is expected from stellar evolution models. The non-rotating Geneva models predict the onset of red supergiants at ~ 4 –4.5 Myr and WC stars at ~ 3.0 –3.5 Myr for solar to twice-solar metallicity (Meynet et al. 1994). Non-zero rotation rates tend to increase both limits by $\sim 20\%$ (Meynet & Maeder 2005). Given the presence of WC stars and absence of red supergiants, we estimate a cluster age of ~ 3.0 –4.5 Myr, assuming coeval formation. Appealing to the Geneva models, we should then expect to still be able to see stars with initial masses ~ 50 –120 $M_{\odot}$ in the cluster. Our age estimate indicates a somewhat lower mass for LBV 1806–20 than given by Eikenberry et al. (2004) since such massive progenitors typically have ages less than 3 Myr (Bond, Arnett, & Carr 1984). A lower mass is also consistent with the claim by Figer et al. (2004) that LBV 1806–20 may be a binary system having $M_{\text{initial}} \sim 130 M_{\odot}$.

The 1806–20 cluster is similar in spectral content to the Quintuplet cluster (Figer et al. 1999), albeit at a much smaller mass. Both have roughly equal numbers of WC and WN stars, although the latter may have twice as many WC’s if the five very red sources for which the Quintuplet was named are indeed dusty WC stars. One minor difference is the presence of one red supergiant in the Quintuplet and none in the 1806–20 cluster. This may be the consequence of low number statistics, as the Quintuplet cluster has four times more massive stars; or it may be an indication that the 1806–20 cluster is slightly younger than the 4 Myr age estimated for the Quintuplet. Assuming 8 stars with $M_{\text{initial}} > 20 M_{\odot}$, we estimate $M_{\text{cluster}} > 1,750 M_{\odot}$, assuming a Salpeter IMF down to 0.8 $M_{\odot}$ (Salpeter 1955). If WR and LBV stars have $M_{\text{initial}} > 50 M_{\odot}$, then $M_{\text{cluster}} > 6,600 M_{\odot}$. The supernova producing SGR 1806–20 likely could not

have triggered formation of the cluster (Eikenberry et al. 2001), because the magnetar is thought to be $<10^4$ yrs old, while the cluster stars are ~ 4 Myr old.

A further important conclusion is that the masses and ages of the cluster stars are consistent with the hypothesis that SGR 1806–20 represents the end state of a particularly massive cluster member. Because more massive stars reach their end states more quickly than less massive stars, the likely mass of the SGR progenitor must be greater than presently observed masses. Assuming that the presently observed stars had $M_{\text{initial}} \gtrsim 50 M_{\odot}$, SGR 1806–20 had a larger initial mass, as suggested by Kaplan et al. (2002). SGR 1900+14 and SGR 0526–66 are also associated with massive stellar clusters (Vrba et al. 2000; Klose et al. 2004), suggesting that magnetars may be short-lived descendants of massive stars. These three

cases support the theoretical prediction that very massive stars lose most of their mass and collapse as neutron stars, not black holes (Fryer & Kalogera 2001; Woosley, Heger, & Weaver 2002). They may also support a model in which magnetars are preferentially produced by stars with large initial mass (Eikenberry et al. 2004).

We acknowledge useful discussions with Jeff Valenti, Andy Fruchter, and Stephan Eikenberry. We thank Keck staff members Randy Campbell and Grant Hill for assistance with the observations. F. N. acknowledges grants AYA2003-02785-E and ESP2002-01627. Finally, we acknowledge a similar result, by independent means, in Gaensler et al. (2005).

REFERENCES

- Bond, J. R., Arnett, W. D., & Carr, B. J. 1984, *ApJ*, 280, 825
- Eikenberry, S. S., Garske, M. A., Hu, D., Jackson, M. A., Patel, S. G., Barry, D. J., Colonna, M. R., & Houck, J. R. 2001, *ApJ*, 563, L133
- Eikenberry, S. S. et al. 2004, *ApJ*, 616, 506
- Feitzinger, J. V., Schlosser, W., Schmidt-Kaler, T., & Winkler, C. 1980, *A&A*, 84, 50
- Figer, D. F., Najarro, F., & Kudritzki, R. P. 2004, *ApJ*, 610, L109
- Figer, D. F., McLean, I. S., & Najarro, F. 1997, *ApJ*, 486, 420
- Figer, D. F. & Kim, S. S. 2002, *ASP Conf. Ser.* 263: *Stellar Collisions, Mergers and their Consequences*, 287
- Figer, D. F., McLean, I. S., & Morris, M. 1999, *ApJ*, 514, 202
- Figer, D. F., Najarro, F., Morris, M., McLean, I. S., Geballe, T. R., Ghez, A. M., & Langer, N. 1998, *ApJ*, 506, 384
- Fryer, C. L. & Kalogera, V. 2001, *ApJ*, 554, 548
- Fuchs, Y., Mirabel, F., Chaty, S., Claret, A., Cesarsky, C. J., & Cesarsky, D. A. 1999, *A&A*, 350, 891
- Gaensler, B. M., McClure-Griffiths, N. M., Oey, M. S., Haverkorn, M., Dickey, J. M., & Green, A. J. 2005, *ArXiv Astrophysics e-prints*, astro-ph/0501563
- Geballe, T. R., Figer, D. F., & Najarro, F. 2000, *ApJ*, 530, 97
- Genzel, R., Thatte, N., Krabbe, A., Kroker, H., & Tacconi-Garman, L. E. 1996, *ApJ*, 472, 153
- Götz, D., Mereghetti, S., Mirabel, I. F., & Hurley, K. 2004, *A&A*, 417, L45
- Hanson, M. M., Conti, P. S., & Rieke, M. J. 1996, *ApJS*, 107, 281
- Hurley, K., Kouveliotou, C., Cline, T., Mazets, E., Golenetskii, S., Frederiks, D. D., & van Paradijs, J. 1999, *ApJ*, 523, L37
- Israel, G. L., Mignani, R., Covino, S., Campana, S., Stella, L., Rea, N., Testa, V., & Marconi, G. 2004, *GRB Circular Network*, 2609, 1
- Kaplan, D. L., Fox, D. W., Kulkarni, S. R., Gotthelf, E. V., Vasisht, G., & Frail, D. A. 2002, *ApJ*, 564, 935
- Klose, S., et al. 2004, *ApJ*, 609, L13
- Koornneef, J. 1983, *A&A*, 128, 84
- Kouveliotou, C., Kippen, M., Woods, P., Richardson, G., Connaughton, V., & McCollough, M. 1998, *IAU Circ.*, 6944, 2
- Kouveliotou, C., Klose, S., Wachter, S., Woods, P., Patel, S., Greiner, J., Stecklum, B., & van der Klis, M. 2004, *GRB Circular Network*, 2607, 1
- Kulkarni, S. R., Matthews, K., Neugebauer, G., Reid, I. N., van Kerkwijk, M. H., & Vasisht, G. 1995, *ApJ*, 440, L61
- LaVine, J. L., Eikenberry, S., & Davis, J. D. 2003, *American Astronomical Society Meeting*, 203,
- Mazets, E. P., Golenetskij, S. V., & Guryan, Y. A. 1979, *Soviet Astronomy Letters*, 5, 343
- Mazets, E. P., Golenetskii, S. V., Ilinskii, V. N., Aptekar, R. L., & Guryan, I. A. 1979, *Nature*, 282, 587
- McLean et al. 1998, *SPIE Vol.* 3354, 566
- Meynet, G., Maeder, A., Schaller, G., Schaerer, D., & Charbonnel, C. 1994, *A&AS*, 103, 97
- Meynet, G., & Maeder, A. 2005, *A&A*, 429, 581
- Ramsay Howat, S. K., et al. 1998, *Proc. SPIE*, 3354, 456
- Salpeter, E. E. 1955, *ApJ*, 121, 161
- Skrutskie, M. F., et al. 1997, *ASSL Vol.* 210: *The Impact of Large Scale Near-IR Sky Surveys*, 25
- Vacca, W. D., Garmany, C. D., & Shull, J. M. 1996, *ApJ*, 460, 914
- van Kerkwijk, M. H., Kulkarni, S. R., Matthews, K., & Neugebauer, G. 1995, *ApJ*, 444, L33
- Vrba, F. J., Henden, A. A., Luginbuhl, C. B., Guetter, H. H., Hartmann, D. H., & Klose, S. 2000, *ApJ*, 533, L17
- Woosley, S. E., Heger, A., & Weaver, T. A. 2002, *Reviews of Modern Physics*, 74, 1015